# Bordetella pertussis/parapertussis Qualitative Real-time PCR



Bordetella pertussis is a gram-negative coccobacillus that causes whooping cough. It occurs worldwide with a high incidence in unvaccinated infants and recently surfaced even in highly vaccinated populations. Symptoms are characterized by violent coughing followed by a high-pitched inspiratory inhalation, or whoop. In non-immunized populations, especially those with malnutrition and a prevalence of other respiratory disease, pertussis is among the most lethal diseases of infants and young children.

Pneumonia is the most common cause of death. The illness caused by B. parapertussis is usually milder than that caused by B. pertussis, although severe cases have been reported. Following the increase of pertussis vaccination coverage, a relative increase of *B. parapertussis* cases have been observed in clinical practice. In vaccinated children, B. pertussis disease frequently presents as a mild disease, clinically difficult to distinguish from diseases associated with coughing caused by B. parapertussis and other bacterial or viral infections. Real-time PCR provides a rapid and sensitive method to determine the presence of target-specific amplifiable nucleic acids in all samples intended for PCR<sup>1-2</sup>. This assay detects both species and distinguishes between them; results include both species. For more information, call the lab at 513-636-9820.

# **Reporting Units:**

Positive/Negative

# Unacceptable Specimens:

- Frozen whole blood
- · Swabs in gel or charcoal media

## **Shipping Conditions:**

- Ambient if sent within 24 hours
- On cold packs if sent >24 hours after collection

## **Testing Schedule:**

Testing for *B. pertussis/parapertussis* is performed every day on first shift. For testing outside of this schedule, call the lab at 513-636-9820. **TAT:** 1-2 days

#### **EPIC Test Code:**

5804023

#### **CPT Code:**

87798 x2

#### **Contact Information:**

Cincinnati Children's Division of Pathology Molecular and Genomic Pathology Services (MGPS)

Phone: 513-636-9820 Fax: 513-517-7099

Email: pathology@cchmc.org

Website: cincinnatichildrens.org/pathology For pricing or billing questions, call 513-636-4261.

## **Shipping Address:**

Cincinnati Children's Hospital Medical Center

Attn: Molecular and Genomic Pathology Services (MGPS) 3333 Burnet Ave, R2.001 Cincinnati. OH 45229

#### References:

- Zuckerman A, Banatvala J, Pattison J, eds. <u>Principles and Practice of Clinical Virology</u>. 2<sup>nd</sup> ed. John Wiley and Sons, West Sussex, United Kingdom. 1990.
- Salamon D, Cummins C, Jones R, et al. Use of multiple in-house and commercial PCR assays for ongoing validation of PCR detection of B. pertussis and B. parapertussis infection. Unpublished data. Columbus Children's Hospital, Department of Laboratory Medicine, Columbus, Ohio. 2009.

Sample Type	Volume Needed	Collection Container
Aspirate: endotracheal tube, tracheal	1mL	Sterile Container
Bronchoalveolar Lavage (BAL) fluid	1mL	Sterile Container
Body Fluids (i.e. pericardial, pleural)*	1mL	Sterile Container
Swab**: nasal, nasopharyngeal, throat	n/a	Red or Green Culturette Swab
Tissue (lung)***	0.3 g	Sterile Container

<sup>\*</sup> Not preferred specimen types for analyzing respiratory agents.

Clinical Lab Index:

Pert: https://www.testmenu.com/cincinnatichildrens/Tests/662803

<sup>\*\*</sup> Red or green top culturette swabs preferred; viral transport media acceptable.

<sup>\*\*\*</sup> Wrap tissue in gauze wetted slightly with sterile saline to keep moist during transport.