

Proteinuria

FAST FACTS

1–5%

of children may be found to have 1+ or greater degree of proteinuria

~10%

of children with ≥ 1 degree of proteinuria will have persistent abnormality after 6 months

less than 2%

of those children will have significant underlying renal disease

In children, protein in the urine is common and usually benign. The most widely used method of screening is the urine dipstick test. Proteinuria is present when urine protein excretion $>4\text{mg}/\text{m}^2/\text{hour}$ or $100\text{mg}/\text{m}^2/\text{day}$.

Spot urine (ideally first morning urine sample) results indicate proteinuria when:

- $>0.2\text{mg protein}/\text{mg creatinine}$ for patients >2 years of age
- $>0.5\text{mg protein}/\text{mg creatinine}$ in patients 6 to 24 months of age.

ASSESSMENT

Perform detailed history focused on description and timing of abdominal/scrotal/leg swelling. Perform complete physical exam focused to evaluate swelling of the abdomen, genitalia and lower extremities. Medical imaging (ultrasound) is unnecessary.

HPE (HISTORY AND PHYSICAL EXAM) RED FLAGS

History of Present Illness:

- Swelling around eyes in the morning
- Swelling in legs in the afternoon, socks leaving prints on legs
- Swollen joints
- Abdominal pain
- High blood pressure: headaches, chest pain, shortness of breath
- Changes in urine output, dysuria
- Skin lesions

Patient History

- Growth history
- Medication intake (NSAIDs, lithium, heavy metals, opioid use particularly heroin)

Family History

- Kidney disease
- Dialysis
- Kidney transplant
- Deafness
- Visual disorders

MANAGEMENT/TREATMENT

If dipstick shows proteinuria, obtain a first morning urine for protein and creatinine ratio.

If urine dipstick is obtained at the time of intercurrent illness and positive for protein, repeat when patient has returned to baseline.

WHEN TO REFER

Refer patients with any of the following to Cincinnati Children's Nephrology:

- Protein/creatinine ratio of >0.2
- Presence of hematuria in addition to proteinuria
- Elevated blood pressure
- Presence of edema and/or rash
- Red flags as described

If you have clinical questions about a patient who with proteinuria, call 513-636-4531 or email nephrology@cchmc.org.

If you would like additional copies of this tool, or would like more information, please contact the Physician Outreach and Engagement team at Cincinnati Children's.

Proteinuria

Patient Presents

Standard Workup

- History of Present Illness
- Family History
- Physical Exam
 - Assess for edema
 - Check blood pressure

HPE (HISTORY AND PHYSICAL EXAM) RED FLAGS

History of Present Illness

- Swelling around eyes in the morning
- Swelling in legs in the afternoon, socks leaving prints on legs
- Swollen joints
- Abdominal pain
- High blood pressure: headaches, chest pain, shortness of breath
- Changes in urine output, dysuria
- Skin lesions


Patient History

- Growth history
- Medication intake (NSAIDs, lithium, heavy metals, opioid use particularly heroin)

Family History

- Kidney disease
- Dialysis
- Kidney transplant
- Deafness
- Visual disorders

Evaluation of Persistent Proteinuria in Children/Adolescents



Method	Indications	Normal Range	Comments
Dipstick testing	Routine screening for proteinuria performed in the office	Negative or a trace in a concentrated urine specimen (specific gravity >1.020) or very concentrated (specific gravity >1.025)	False positive can occur if urine is very alkaline (pH >8.0)
24-hour urine for proteinuria and creatinine excretion	Quantitation of proteinuria as well as creatinine clearances	<100 mg/m ² /24 h	More accurate than spot urine analysis
Spot urine for protein/creatinine ratio, preferably on first morning urine	Semi-quantitative assessment of proteinuria	<0.2 mg protein/mg creatinine in children >2 years old <0.5mg protein/mg creatinine in children age 6–24 months	Simplest method to detect proteinuria. Less accurate than 24-hour test
Micro-albuminuria	Assess risk of progressive glomerulopathy	<30 mg urine albumin/g creatinine on first morning urine	Therapy should be intensified in diabetics with MA in DM

For urgent issues, or to speak with the specialist on call 24/7, call the Physician Priority Link® at 1-888-987-7997.